Economia agro-alimentare / Food Economy

An International Journal on Agricultural and Food Systems

Vol. 27, Iss. 3, Art. 13, pp. 1-24 - ISSNe 1972-4802

DOI: 10.3280/ecag2025oa20804



Grazing Impacts on Biodiversity, Carbon Cycle, Water Efficiency, and Animal Welfare: Review

Giulia Pastorelli^{*,a}, Ilaria Falconi^a, Maria Assunta D'Oronzio^a, Raffaella Pergamo^a

^a CREA Research Centre for Agricultural Policies and Bioeconomy, Italy

Abstract

This review examines the role of pasture-based livestock systems in promoting agri-environmental sustainability across four key dimensions: biodiversity, water-use efficiency, carbon cycle, and animal welfare. Despite growing political and scientific interest in extensive grazing, the existing literature remains fragmented and limits a comprehensive understanding of grazing's multidimensional impacts. To address this gap, we conducted a review of studies published between 2010 and 2025, following the Cochrane Handbook and applying the PICO (Population, Intervention, Comparison, Outcome) framework. The analysis focused exclusively on studies conducted in Europe to explore the state of the art on the topic and analyze future policy implications. The results indicate that moderate and well-managed grazing improves biodiversity, increases water-use efficiency, and promotes soil carbon sequestration, especially through rotation practices and tree integration. Access to pasture improves animal welfare, although outcomes vary depending on infrastructure and environmental conditions. The review emphasizes the need for integrated approaches that combine traditional knowledge, environmental design, and precision tools to maximize the sustainability of grazing systems.

Article info

Type: Review Submitted: 01/08/2025 Accepted: 20/10/2025 Available online: 20/11/2025

JEL codes: N54, O13, P28

Keywords:

Water-use Carbon cycle Animal welfare Biodiversity grazing systems

Managing Editor: Veronica Manganiello

^{*} Corresponding author: Giulia Pastorelli - CREA Research Centre for Agricultural Policies and Bioeconomy, Rome (Italy). E-mail: giulia.pastorelli@crea.gov.it.

Introduction

Agriculture plays a crucial role in safeguarding ecosystem services, food security, and landscape preservation. However, demographic expansion, resource competition, geopolitical tensions, and climate change are placing increasing pressure on agricultural systems. In this context, halting biodiversity loss is a priority for the European Union, as outlined in the Farm to Fork and Biodiversity 2030 Strategies (Fraser *et al.*, 2022). The transdisciplinary One Health approach can strengthen resilience and prevention against environmental risks, antimicrobial resistance (AMR), and climate change (Rodriguez, 2024). Within agri-livestock systems, the debate around production intensification is growing. Extensive farming systems, through the diversification of grazing surfaces and species, promote biodiversity and soil fertility (Delaby *et al.*, 2020).

Despite growing interest in these systems, literature lacks comprehensive data on the environmental impact of livestock farming, as most studies have primarily focused on carbon footprint (Tittonell, 2021; Opio *et al.*, 2013), neglecting other aspects such as biodiversity, land stewardship, energy efficiency, and sociocultural values.

Elorriaga *et al.* (2013) highlight that the concentration of manure in intensive farming systems and the frequent use of antimicrobials result in significant environmental impacts, including water resource pollution with potential negative consequences for human health. Similarly, Jechalke *et al.* (2014) emphasize that manure spreading practices associated with such production systems may pose high contamination risks and promote the development of antimicrobial resistance.

In the present review, we adopt the PICO framework (Population, Intervention, Comparison, Outcome) to define the scope of analysis and guide the construction of search strings. Following Cochrane guidelines (Cumpston *et al.*, 2024), the study is based on the effects of pasture-based grazing systems in relation to four critical dimensions of agro-environmental sustainability: biodiversity, water use efficiency, carbon cycle, and animal welfare.

1. Background

In recent years, consumers have shown growing interest in purchasing products derived from sustainable agricultural systems, which are perceived as more respectful of environmental integrity and animal welfare. This trend reflects a broader societal concern over the ecological footprint of food production and a demand for transparency in how ecosystems are managed and preserved. Ecosystems provide essential services, such as food

provision, water regulation, and climate mitigation, that underpin human wellbeing and economic development. Yet, these contributions are rarely reflected in traditional economic indicators, creating a disconnect between environmental value and policy action (Vargas et al., 2024). Across many regions, especially those undergoing accelerated agricultural intensification. the lack of long-term planning and water resource monitoring has led to significant sustainability challenges, particularly in terms of water use (Barsotti et al., 2022). At the same time, the abandonment of traditional land use practices, common throughout Europe in recent decades, has had profound impacts on agricultural landscapes, resulting biodiversity loss (Ponzetta et al., 2010). Indeed, it is estimated that 75% of grassland habitats in the European Union are in poor or bad condition, with land abandonment and overgrowth representing key threats (Hessle et al., 2023). The integration of food production with biodiversity conservation has thus emerged as a critical challenge for sustainable development (Gallé et al., 2020). This is particularly urgent in the context of the ongoing need for ecological intensification of farming systems, improving agricultural productivity while minimizing environmental degradation and maintaining ecosystem functions (Beye et al., 2019). Future grazing systems are expected to play a central role in this transition by supplying nutrient- dense food within lower-input systems that reduce reliance on fossil fuels and agrochemicals, and enhance environmental. biodiversity, and animal welfare outcomes (Delaby et al., 2020). Lower-input pasture-based systems offer several advantages: they utilize feed sources inedible to humans (e.g., grasslands) rather than human-edible grains (e.g., soya), and they reduce feeding costs and the environmental footprint of production. Additionally, these systems often align with consumer preferences for higher animal welfare and environmentally sustainable meat products. In this context, the preservation and promotion of autochthonous livestock breeds are of strategic importance. These breeds not only embody genetic diversity crucial for adaptation to local environments, but also support rural development, cultural identity, and food and nutrition security, especially in marginal and pasture-based systems (Agradi et al., 2023). According to Devincenzi et al. (2021), grazing behaviour management plays a crucial role in improving animal welfare, product quality, and ecosystem services.

2. Materials and methods

This review was conducted following the methodological principles outlined in the Cochrane Handbook for Systematic Reviews of Interventions, and the review design was structured using the PICO model (Population, Intervention, Comparison, Outcome) (Table 1) (Sala *et al.*, n.d.) to formulate

the research question and organize the thematic domains. The aim was to identify and synthesize scientific literature on the environmental and animal welfare effects of pasture-based grazing systems.

Table 1 - PICO model

P	Population/Problem	Livestock production systems
I	Intervention	Effects of grazing on biodiversity, water use efficiency, carbon sequestration, and animal welfare
C	Comparison	Comparison between livestock systems and farming methods at the European and global level
o	Outcome	Scientific evidence on the most frequently addressed and associated sustainability themes

Source: Authors' elaboration.

To ensure methodological rigor and transparency, inclusion and exclusion criteria were defined a priori, and reinforced by specific guidelines for systematic reviews in the agri-food sector, which emphasize the importance of clearly defining the study question and selecting relevant study designs (Sargeant *et al.*, 2005).

Studies were selected based on their relevance to at least one of the following dimensions: biodiversity, water use efficiency, carbon cycling, and animal welfare in grazing-based livestock systems.

Table 2 - Inclusion and exclusion Criteria

Inclusion Criteria	Exclusion Criteria	
Studies evaluating grazing effects on biodiversity, water use, carbon cycle, or animal welfare	Studies not addressing at least one of the four sustainability dimensions	
Articles, Narrative reviews	commentaries, opinion pieces	
From 2010 to 2025	Published before 2010	
Studies conducted in Europe	Studies not conducted in Europe	
Full-text available	Abstract only or inaccessible full text	
	Studies evaluating grazing effects on biodiversity, water use, carbon cycle, or animal welfare Articles, Narrative reviews From 2010 to 2025 Studies conducted in Europe	


Source: Authors' elaboration.

The initial identification of records was carried out manually via Scopus. Search strings were constructed using combinations of key terms related to the four thematic categories:

- Pasture grazing and biodiversity.
- Pasture grazing and water use.
- Pasture grazing and carbon farming.
- Pasture grazing and animal welfare.

The flow diagram representing the systematic selection process for all included studies follows the PRISMA (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) method (Page *et al.*, 2021). Preliminary exclusion criteria were applied directly on Scopus through filters, including publication date (from 2010 to 2025), availability of full text (open access), and geographic context (with a focus on European studies).

Figure 1 - PRISMA-based Schema



Source: Author's elaboration on base Page M. J. et al. BMJ 2021.

The next stage (reports sought for retrieval, n = 105) involved retrieval of all texts and a more careful assessment of methodological appropriateness

and thematic relevance. It is worth noting that a small number of duplicate records were identified at this stage (n = 3). This limited overlap was due to the use of a single database (Scopus) and four separate thematic searches (via keywords). Separation by domain initially prevented automatic detection of duplicates, which was resolved during eligibility screening.

3. Results

3.1. Literature distribution

Data extracted from the selected studies were synthesised using a narrative approach, given the heterogeneity of study designs, geographical contexts, and methodological approaches. The synthesis was structured thematically along the four main dimensions defined in the research framework: (1) biodiversity, (2) water use efficiency, (3) carbon cycle, and (4) animal welfare. Each study was categorized based on its thematic focus and geographical location, selecting only European studies.

A total of 765 studies were initially retrieved from Scopus after keyword filtering. The thematic distribution highlights a predominant focus on water use (49.4%), followed by biodiversity (29.5%), animal welfare (12.6%), and carbon (8.5%). A temporal analysis reveals a shift in research interest: while the link between environmental dimensions such as grazing and biodiversity or water use has been explored for over a decade, most studies on carbon sequestration and animal welfare are more recent, with over 90% of articles in these categories published after 2010 (Table 3).

Table 3 - Temporal distribution of literature

Thematic Area	No. of Articles	% of Total	Articles 2010-2025	% of Category
Pasture grazing & biodiversity	227	29.5%	184	81%
Pasture grazing & water use	377	49.4%	265	70%
Pasture grazing & carbon farming	65	8.5%	63	96%
Pasture grazing & animal welfare	96	12.6%	88	92%
Total	765	100%	598	_

Source: Author's elaboration.

This suggests a growing awareness of the role of pasture-based livestock systems in climate change mitigation and the sustainability of livestock farming.

Another dimension explored was the geographical distribution of the literature (Table 4). Of the 764 studies, only 260 (34%) focused on European countries, revealing a significant knowledge gap related to the four explored dimensions of grazing, particularly in relation to water use and carbon sequestration. Thematic areas such as animal welfare (52.1%) and biodiversity (54.2%) have a relatively larger share of EU-based studies, while water use lags significantly behind (18.3%), despite its relevance in a context of increasing climate variability and water stress in southern Europe.

Table 4 - Geographical distribution of the literature

Thematic Area	No. of Articles	EU Articles	% EU of Total
Pasture grazing & biodiversity	227	123	54.2%
Pasture grazing & water use	377	69	18.3%
Pasture grazing & carbon farming	65	18	27.7%
Pasture grazing & animal welfare	96	50	52.1%
Total	765	260	34.0%

Source: Author's elaboration.

The preliminary evidence emerging from this review highlights the evolving role of grazing systems across multiple sustainability dimensions in the European context. While the environmental role of pasture-based livestock farming has long been acknowledged in terms of biodiversity and water dynamics, emerging challenges such as climate change mitigation, greenhouse gas emissions, and ethical livestock management are gaining increasing attention. However, the geographic and thematic imbalance in the literature suggests a need for greater integration between sustainability dimensions and for enhanced focus on European systems, particularly in water governance and carbon-related practices.

3.2. *Narrative synthesis*

The following narrative synthesis presents the key findings emerging from the selected studies, organized according to the four sustainability dimensions examined in this review. Given the heterogeneity of study designs, geographic contexts, and methodological approaches, results are discussed thematically to highlight recurring patterns, relevant contrasts, and emerging insights. The synthesis focuses on how pasture-based livestock system have been studied in relation to each dimension, offering a comprehensive overview of their environmental and animal welfare implications across different European contexts.

3.2.1. Grazing and biodiversity

Pasture grazing plays a central role in the conservation of biodiversity across European landscapes. Numerous studies confirm that grazing, when properly managed, enhances both plant and habitat diversity, especially in semi-natural and marginal environments.

Moderate grazing intensity, for instance, has been associated with increased plant and pollinator richness (Freschi et al., 2015), and with improved habitat heterogeneity in low- input pastures (Tonn et al., 2019). In Mediterranean systems, Candel-Pérez et al. (2024) and Fracchiolla et al. (2017) demonstrate how intensive or unmanaged grazing reduces plant richness, favouring disturbance-tolerant species. By contrast, targeted restoration interventions such as mowing or reseeding (Ponzetta et al., 2010; Sitzia et al., 2010) have proven effective in reversing biodiversity loss in abandoned grasslands. Similar dynamics emerge in Polish meadows, where grazing influenced species abundance and beetle community composition (Szyszko-Podgórska et al., 2024; Czarnecki et al., 2015), and in Sweden, where the reintroduction of grazing restored functional invertebrate assemblages (Steiner et al., 2016).

Animal biodiversity and habitat structure also benefit from appropriate grazing patterns. Schmitz & Isselstein (2020) and van Klink *et al.* (2016) show that grazing regime, more than species, determines structural diversity and ecosystem function. Dumont *et al.* (2020) found that pasture size altered vegetation patchiness, while Fleurance *et al.* (2016) noted positive insect responses to moderate stocking rates. Furthermore, Wróbel *et al.* (2020) highlight how moderate sheep grazing sustains both plant richness and pastoral value in mountain ecosystems. Landscape-level perspectives reinforce these findings. Plieninger *et al.* (2015) identify wood-pastures as biodiversity hotspots tied to cultural values. In a broader EU context, climate

change adds additional complexity Dibari *et al.* (2015) report reduced pasture suitability in the Apennines, threatening grazing-dependent biodiversity. Similarly, Lennartsson & Westin (2023) document how the loss of traditional seasonal grazing affects grassland resilience. At policy level, Roeder *et al.* (2010) and Asheim *et al.* (2020) argue for improved incentives to support pasture-based systems, while Phelps & Kaplan (2017) call for better land-use frameworks integrating biodiversity goals.

Finally, the role of local breeds and traditional practices emerges clearly. Studies from Agradi *et al.* (2023), and Verduna *et al.* (2020) link transhumance and autochthonous livestock to greater landscape heterogeneity and ecological sustainability. Potenza & Fedele (2011) emphasize that well-managed grazing, aligned with agri-environmental rules (e.g. GAEC) of Common Agriculture Policy (CAP), supports floristic richness even in marginal terrains.

3.2.2. Grazing and water use

The management of grazing systems plays a pivotal role in shaping water use efficiency (WUE), with direct implications for soil conservation, hydrological dynamics, and reduced irrigation demand in livestock production. Several studies confirm that that well-managed grazing, such rational grazing, particularly in semi-arid regions, can optimize water use without compromising animal productivity. For instance, Ates *et al.* (2013) showed that deficit irrigation (50–75%) in pasture-based sheep systems improved WUE while maintaining both yield and animal performance.

Beyond water requirements, grazing significantly influences soil structure and hydrological functioning. Moret-Fernández et al. (2011) reported that grazing alters the hydro-physical properties of soils, notably reducing infiltration rates and increasing bulk density, particularly in non-gypsiferous substrates. Similarly, Minea et al. (2022), observed that small ruminants' grazing intensified surface runoff, reduced infiltration, and increased erosion in hilly and mountainous terrains, ultimately affecting local water balances. Despite such potential risks, appropriately managed grazing can promote hydrological stability and improve resource efficiency. In semi-arid Mediterranean pastures characterized by chalky soils, Moret-Fernández et al. (2021) demonstrated that intensive livestock grazing significantly reduced surface water infiltration due to trampling-induced crusting, while having minimal impact on soil matrix water retention. In contrast, moderate grazing intensity did not negatively affect soil hydraulic properties, indicating its potential suitability as a sustainable management strategy for soil conservation under these specific drought conditions.

From a livestock behavior standpoint, Tsiobani et al. (2023) observed that water availability influences buffalo spatial distribution in Mediterranean pasture systems, as animals tend to cluster near water points, an effect with notable implications for grazing patterns and landscape pressure. Pasturebased systems also contribute to reducing overall water consumption in meat production. According to Dillon et al. (2018), in recent decades, permanent grasslands in Europe have declined by 14%, leading to an increased reliance on irrigated crops for livestock feed. Comparative analyses of production systems indicate that water use to produce 1 kg of beef is 30-50% lower in grazing-based systems compared to intensive systems. Furthermore, grazing contributes to improved soil moisture and reduced evaporation, thereby decreasing the need for artificial irrigation. In support of this, modeling by Simionesei et al. (2018) on Portuguese montado systems confirmed that irrigation can increase biomass yield, but also raised transpiration, suggesting that grazing management needs to balance productivity with water use. Grazed pasture, being the lowest-cost and rain-fed feed source in temperate Europe, enhances the self-sufficiency and environmental sustainability of dairy systems by reducing reliance on externally irrigated feed crops and buffering against water-related constraints (Hennessy et al., 2020). Finally, Bernués et al. (2011) emphasized that sustainable grazing practices contribute to water security by reducing irrigation needs and improving the resilience of Mediterranean farming systems. Extending the grazing season from 252 to 300 days has been shown to decrease the need for irrigation water and improve the forage self-sufficiency of livestock farms.

3.2.3. Grazing and Carbon Cycle

A growing body of empirical and theoretical research demonstrates that well-managed pasture-based systems play a significant role in enhancing carbon sequestration and mitigating greenhouse gas (GHG) emissions in a variety of agroecological contexts. Tölgyesi *et al.* (2023) highlight the multifunctional benefits of sparse trees in temperate tree pastures in Hungary and Romania, where they achieved increased carbon stocks and biodiversity without compromising livestock productivity. These findings are consistent with those of Smetanová *et al.* (2019), who report that traditional extensive grazing in the Czech Republic maintains higher soil carbon levels than abandoned lands, resulting in benefits for biodiversity and landscape conservation. Complementary findings from Poland show that rotational grazing of deer improves forage quality and alters plant chemical composition (Kulik *et al.*, 2023), while also modulating soil chemical and enzymatic properties (Futa *et al.*, 2024), albeit with a reduction in total organic carbon

during the winter. Studies focusing on carbon footprints also point to the benefits of grazing systems. Morais *et al.* (2018) report lower greenhouse gas emissions per kg of product in extensive grazing systems using local breeds, from the Azores showing a carbon footprint of 0.83 kg CO₂e/kg of milk, among the lowest in Europe. Morais *et al.* (2025) further highlight the variability of GHG emissions (15-124 kg CO₂e/100 g protein), noting that pasture-based systems consistently outperform in terms of emissions reductions. Hessle and Danielsson (2024) estimate that the restoration of semi-natural grasslands in Sweden can be achieved through favourable conservation status, strategic reductions in livestock density rates, and partial mowing. Carswell *et al.* (2024) report that reseeding pastures with legumerich swards improves nitrogen use efficiency in beef and sheep systems, without significantly increasing productivity, while also enhancing carbon sequestration in established swards.

O'Brien et al. (2016) provide life-cycle assessment evidence that moderate intensification improves carbon efficiency in Irish sheep systems, although the results are highly dependent on assumptions regarding soil carbon sequestration. At the system design level, Barron et al. (2021) propose an integrated assessment framework for pasture-based dairy sheep farming systems in Spain, identifying soil carbon sequestration as a key sustainability indicator. Meanwhile, Pauler et al. (2023) demonstrate that carbon storage in mountain pastures is influenced by topography and climate (e.g., slope, temperature), with higher forage productivity correlating with greater carbon storage but reduced biodiversity, highlighting key ecosystem trade-offs. Recent work by Zhu et al. (2023) also confirms that high-resolution satellite data combined with eddy covariance flux towers can reliably extrapolate carbon fluxes at the field scale in managed European grazing systems, offering promising tools to monitor and assess pasture-based climate mitigation strategies.

3.2.4. Grazing and Animal Welfare

Hesselmann *et al.* (2025) demonstrate that allowing dairy cows to graze on multispecies swards supports natural foraging behavior and individual dietary self-selection, enhancing welfare through behavioral autonomy. Similarly, Molle *et al.* (2022) emphasize the role of planned rotational grazing (PTG) in supporting welfare by improving mobility, milk quality, and promoting natural behaviors. The traditional knowledge of shepherds, as documented by Rivera *et al.* (2022), also contributes to welfare, with the use of over 200 medicinal plants for animal care in transhumant systems. The influence of grazing intensity and management was assessed by Crossley *et al.* (2022),

who reported only minor welfare differences between treatment groups, with better locomotor conditions and comfort observed in cows with longer grazing periods. Likewise, Crump et al. (2019) found that overnight pasture access improved cow welfare by enhancing lying time, activity, and behavioral synchrony. However, temporary indoor confinement, as shown by Enriquez-Hidalgo et al. (2018), negatively affected lying behavior and locomotion, especially in Jersey cows, while O'Driscoll et al. (2019) observed reduced lying behavior due to limited pasture allowance. Environmental enrichment and infrastructure also play a key role. Delaby et al. (2020) highlight how hedgerows and wooded areas provide shade and shelter. improving thermal comfort and reducing disease risk. Turini et al. (2025) confirms this by showing that access to water and shade during hot periods mitigates haemoconcentration and metabolic stress in sheep, compared to animals exposed to harsher conditions. Technological solutions were explored by Marchegiani et al. (2025), who report that Precision Livestock Farming (PLF) tools enhance welfare via real-time health monitoring and optimized grazing management, although adoption remains limited. Similarly, Bosona & Gebresenbet (2021) suggest that mobile automated milking systems can promote cow-calf contact and reduce stress in pasture-based setups. McCormick & Stokes (2023) note that GPS collars, if well-managed, offer benefits in extensive systems despite potential welfare concerns related to invisible or fluctuating boundaries. Behavior-based metrics are gaining ground but still face challenges. Aubé et al. (2023) found that avoidance distance at pasture (ADP) is not yet a reliable individual-level indicator for human-animal relationship assessment. O'Leary et al. (2020) observed that mild and moderate lameness in cows did not consistently correlate with behaviour patterns, suggesting limitations in current accelerometerybased detection. Research in other species reinforces the welfare potential of pasture-based systems. Battini et al. (2021) adapted the AWIN (Welfare Indicators Network) protocol for goats and identified generally good welfare, despite concerns about body condition and water access. Agradi et al. (2023) and Niero et al. (2022) confirm that well-managed alpine and summer pastures support animal health and milk quality without negative impacts on udder health. Welfare implications of landscape use were examined by Larraz et al. (2024) who used GPS data to analyze habitat preferences of free-grazing sheep, suggesting better alignment of resource distribution with animal behavior. In a similar ecological context, Rutherford et al. (2021) highlight that incorporating surplus male dairy calves into pasture-based beef production systems represents a more welfare-friendly alternative to practices such as long- distance export. Strand et al. (2019) highlight a concret concern in outfield systems, predation by large carnivores increases stress, injury, and mortality among sheep, suggesting that welfare benefits

of grazing systems are context dependent. Grodkowski *et al.* (2023) further emphasize that organic certification alone does not always guarantee superior welfare, especially when compared to well-managed conventional systems.

Finally, Ripamonti *et al.* (2024) report no significant welfare differences between cattle raised in silvopastoral and open pasture systems, as evidenced by stable cortisol levels. Avondo *et al.* (2013) link pasture-based feeding to enriched milk and cheese quality, with bioactive compounds that not only enhance product value but may also support animal health.

4. Discussion

While this review focused on the environmental and animal welfare impacts of pasture-based livestock systems in Europe, it is essential to contextualize the findings within a broader sustainability framework, including economic viability and global perspectives.


Evidence from non-European studies, in line with this review, highlights substantial variability in sustainability outcomes even among comparable grazing systems. As Broom (2021) argues, the mere classification of a system as "extensive" or "pasture-based" does not ensure sustainability; rather, outcomes depend on context-specific factors, including grazing pressure, animal density, pasture design, and management strategies. For instance, the same rotational grazing model may lead to biodiversity gains in one context and neutral or even negative effects in another, depending on how it is implemented (Ripoll-Bosch *et al.*, 2013). This calls for adaptive and locally informed grazing strategies rather than blanket recommendations.

Moreover, although our review restricted data inclusion to European studies, relevant global literature offers valuable insights that reinforce and expand upon our conclusions. For example, silvopastoral systems in Latin America have demonstrated that integrating tree cover into grazing landscapes can yield positive outcomes for biodiversity, productivity, and animal welfare (Broom *et al.*, 2013; Broom, 2017). These systems offer functionally similar models to Mediterranean agroforestry and could inspire design adaptations in Europe's southern regions. Additionally, Broom (2022) proposes the use of sustainability scoring to assess the balance of trade-offs across environmental, animal, and economic dimensions a tool potentially adaptable to European contexts.

Crucially, the economic sustainability of grazing systems must be further emphasized. Although often overlooked in environmental reviews, financial viability is essential for long-term adoption by farmers. Studies by Bernués *et al.* (2014) highlight that pasture-based systems contribute to rural livelihoods and resilience, especially in marginal areas where alternative agricultural

models are less feasible. These systems also tend to show lower dependence on external inputs and greater resilience to market volatility, due to self-sufficiency in forage and reduced feed costs (Toro-Mujica *et al.*, 2015; Ripoll-Bosch *et al.*, 2013). Furthermore, as FAO (Chará *et al.*, 2019) notes, despite lower short-term productivity, these systems offer robust long-term benefits for food security and socio-economic stability under climate stress.

Figure 2 - Factors influencing the sustainability of extensive or pasture-based system



Source: Author's elaboration.

Conclusions

This review substantially confirms the contribution of well-managed grazing systems to global agro-environmental sustainability. Properly managed livestock farming systems based on pasture can offer significant benefits across all key sustainability dimensions: environmental, economic, climatic and animal welfare. The analysed scientific evidence indicates that practices such as rotational grazing, holistic management, agro-livestock integration, and adapting grazing practices to local ecological and climatic conditions can improve water use efficiency (WUE), optimise hydrological

dynamics, increase organic carbon sequestration in the soil, and preserve biodiversity without compromising livestock performance. In particular, adopting strategies such as deficit irrigation in pastoral systems in semiarid regions has been shown to significantly reduce water consumption while maintaining comparable yields, thereby contributing to the ecological resilience of these areas. Conversely, the structure and management of pastures have a profound influence on the physical, chemical and enzymatic properties of the soil, thus confirming the interdependence between livestock practices, soil health and environmental quality. Furthermore, recent research highlights the importance of breed-specific grazing behaviour. grazing time distribution and vegetative structure in determining animal performance and the overall sustainability of systems. Another important factor is animal welfare. Extensive grazing encourages natural behaviour and improves feeding autonomy, promoting better physical and behavioural health in line with the principles of the One Health framework. In particular, in low-input farming, animals play a central role as efficient converters of non- edible plant biomass into animal-based food. For instance, dairy cows transform grass into milk, supporting the circular use of resources and contributing to food security. Increasing the proportion of grazed forage in the annual feed plan is associated with lower production costs, higher profitability, and greater farm resilience. However, these benefits do not occur automatically and depend on several factors, including grazing intensity, infrastructure quality, soil and climate conditions, landscape, and political and institutional support. Integrating traditional knowledge with advanced technological tools, such as sensors, GPS tracking, decision support systems and precision farming technologies, offers concrete prospects for adaptive pasture management in response to climate change and increasing environmental variability. These emerging tools offer promising opportunities to improve the monitoring and optimisation of interactions between animals, vegetation and soil. However, there are still obstacles to adoption and specific challenges, including water scarcity, predation risks and land fragmentation, which require contextualised, multisectoral solutions. In light of this evidence, it is clear that extensive livestock systems can represent a strategic ecological safeguard for landscape regeneration, soil fertility improvement and climate change mitigation, rather than being a threat to the environment. Future agricultural development strategies should therefore recognise the functional role of grazing within an integrated ecosystem approach, promoting targeted support policies and investments in transdisciplinary research, as well as coordinated actions at local, national and global levels. Only through such a systemic, multi-level vision will it be possible to exploit the full regenerative potential of pastoral systems and contribute concretely to achieving global climate, environmental and socio-economic objectives.

References

- Agradi, S., Munga, A., Barbato, O., Palme, R., Tarhan, D., Bilgiç, B., Dokuzeylül, B., Ercan, A. M., Or, M. E., Brecchia, G., Curone, G., Draghi, S., Vigo, D., Marongiu, M. L., González-Cabrera, M., & Menchetti, L. (2023). Goat hair as a bioindicator of environmental contaminants and adrenal activation during vertical transhumance. *Frontiers in Veterinary Science*, 10. Doi: 10.3389/fvets.2023.1274081.
- Asheim, L. J., Thorvaldsen, P., & Rivedal, S. (2020). Policy measures to preserve Norwegian coastal and fjord landscapes in small-scale farming systems. *Environmental Science and Policy*, 104, 43-51. Doi: 10.1016/j.envsci.2019.10.017.
- Ates, S., Isik, S., Keles, G., Aktas, A. H., Louhaichi, M., & Nangia, V. (2013). Evaluation of deficit irrigation for efficient sheep production from permanent sown pastures in a dry continental climate. *Agricultural Water Management*, 119, 135-143. Doi: 10.1016/j.agwat.2012.12.017.
- Aubé, L., Mollaret, E., Mialon, M. M., Mounier, L., Veissier, I., & de Boyer des Roches, A. (2023). Measuring the human-animal relationship in cows by avoidance distance at pasture. *Applied Animal Behaviour Science*, 265. Doi: 10.1016/j.applanim.2023.105999.
- Avondo, M., Secchiari, P., Battaglini, L. M., Bonanno, A., & Pulina, G. (2013). Soil, pasture and animal product quality. *Italian Journal of Agronomy*, 8(3), 141-148. Doi: 10.4081/ija. 2013.e19.
- Barron, L. J. R., Andonegi, A., Gamboa, G., Garmendia, E., García, O., Aldai, N., & Aldezabal, A. (2021). Sustainability assessment of pasture-based dairy sheep systems: A multidisciplinary and multiscale approach. Sustainability (Switzerland), 13(7). Doi: 10.3390/su13073994.
- Barsotti, M. P., de Almeida, R. G., Macedo, M. C. M., Laura, V. A., Alves, F. v., Werner, J., & Dickhoefer, U. (2022). Assessing the freshwater fluxes related to beef cattle production: A comparison of integrated crop-livestock systems and a conventional grazing system. *Agricultural Water Management*, 269. Doi: 10.1016/j.agwat.2022.107665.
- Battini, M., Renna, M., Giammarino, M., Battaglini, L., & Mattiello, S. (2021). Feasibility and Reliability of the AWIN Welfare Assessment Protocol for Dairy Goats in Semi-extensive Farming Conditions. *Frontiers in Veterinary Science*, 8. Doi: 10.3389/fvets.2021.731927.
- Bernués, A., Olaizola, R. A. ^c, Villalba, D., & Casasús I. (2011). Sustainability of pasture-based livestock farming systems in the European Mediterranean context: Synergies and trade-offs. *Livestock Science*, *139*(1-2), July, 44-57. Doi: 10.1016/j. livsci.2011.03.018.
- Bernués, A., Rodríguez-Ortega, T., Ripoll-Bosch, R., & Alfnes, F. (2014). Sociocultural and economic valuation of ecosystem services provided by Mediterranean mountain agroecosystems. *PLoS ONE*, *9*(7). Doi: 10.1371/journal.pone.0102479.
- Beye, H., Taube, F., Donath, T. W., Schulz, J., Hasler, M., & Diekötter, T. (2023). Species Enriched Grass-Clover Pastures Show Distinct Carabid Assemblages and Enhance Endangered Species of Carabid Beetles (Coleoptera: Carabidae) Compared to Continuous Maize. *Land*, *12*(4). Doi: 10.3390/land12040736.

- Bosona, T., & Gebresenbet, G. (2021). Multipurpose simulation model for pasture-based mobile Automated Milking and Marketing System, Part-I: Pasture, milk yield, and milk marketing characteristics. *Computers and Electronics in Agriculture*, 190. Doi: 10.1016/j.compag.2021.106212.
- Broom, D. M. (2017). Components of sustainable animal production and the use of silvopastoral systems. *Revista Brasileira de Zootecnia*, 46(8), 683-688. Doi: 10.1590/S1806-92902017000800009.
- Broom, D. M. (2021). A method for assessing sustainability, with beef production as an example. *Biological Reviews*, 96(5), 1836-1853. Doi: 10.1111/brv.12726.
- Broom, D. M. (2022). Animal board invited opinion paper: The use of sustainability scoring to evaluate food production and prepare for the future. *Animal*, *16*(12). Doi: 10.1016/j.animal.2022.100680.
- Broom, D. M., Galindo, F. A., & Murgueitio, E. (2013). Sustainable, efficient livestock production with high biodiversity and good welfare for animals. *Proceedings of the Royal Society B: Biological Sciences*, 280(1771). Doi: 10.1098/rspb.2013.2025.
- Candel-Pérez, D., Lucas-Borja, M. E., Plaza-Álvarez, P. A., Carmona Yáñez, M. D., Soria, R., Ortega, R., Miralles, I., Miralha, L., & Zema, D. A. (2024). Effects of grazing on soil properties in mediterranean forests (Central-Eastern Spain). *Journal of Environmental Management*, 354. Doi: 10.1016/j.jenvman.2024.120316.
- Candel-Pérez, D., Lucas-Borja, M. E., Plaza-Álvarez, P. A., Carmona Yáñez, M. D., Soria, R., Ortega, R., Miralles, I., Miralha, L., & Zema, D. A. (2024). Effects of grazing on soil properties in mediterranean forests (Central-Eastern Spain). *Journal of Environmental Management*, 354. Doi: 10.1016/j.jenyman.2024.120316.
- Carswell, A. M., Gongadze, K., Misselbrook, T. H., & Wu, L. (2019). Impact of transition from permanent pasture to new swards on the nitrogen use efficiency, nitrogen and carbon budgets of beef and sheep production. *Agriculture, Ecosystems and Environment*, 283. Doi: 10.1016/j.agee.2019.106572.
- Chará J., Reyes E., Peri P., Otte J., Arce E., Schneider F. (2019). Silvopastoral Systems and their Contri bution to Improved Resource Use and Sustainable Development Goals: Evidence from Latin America. FAO, CIPAV and Agri Benchmark, Cali, 60 pp. Licence: CC BY-NC-SA 3.0 IGO.
- Crossley, R. E., Bokkers, E. A. M., Browne, N., Sugrue, K., Kennedy, E., Engel, B., & Conneely, M. (2022). Risk factors associated with the welfare of grazing dairy cows in spring-calving, hybrid pasture-based systems. *Preventive Veterinary Medicine*, 204. Doi: 10.1016/j.prevetmed.2022.105640.
- Crump, A., Jenkins, K., Bethell, E. J., Ferris, C. P., & Arnott, G. (2019). Pasture access affects behavioral indicators of wellbeing in dairy cows. *Animals*, *9*(11). Doi: 10.3390/ani9110902.
- Cumpston, M., Flemyng, E., Thomas, J., Higgins, J. P. T., Deeks, J. J., & Clarke, M. J. (2024). Chapter I: Introduction [last updated August 2023]. In: Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch V. A. (eds.), Cochrane Handbook for Systematic Reviews of Interventions version 6.5. Cochrane. Available from training.cochrane.org/handbook.
- Delaby, L., Finn, J. A., Grange, G., & Horan, B. (2020). Pasture-Based Dairy Systems in Temperate Lowlands: Challenges and Opportunities for the Future.

- In Frontiers in Sustainable Food Systems (Vol. 4). Frontiers Media S.A. Doi: 10.3389/fsufs.2020.543587.
- Devincenzi, T., Coppa, M., & Cabiddu, A. (2021). Editorial: Ruminant Grazing Behavior: A Tool to Improve Product Quality and Ecosystem Services. In *Frontiers in Veterinary Science* (Vol. 8). Frontiers Media S.A. Doi: 10.3389/fvets.2021.744200.
- Dibari, C., Argenti, G., Catolfi, F., Moriondo, M., Staglianò, N., & Bindi, M. (2015). Pastoral suitability driven by future climate change along the apennines. *Italian Journal of Agronomy*, 10(3), 109-116. Doi: 10.4081/ija.2015.659.
- Dillon P.G. (2018). The evolution of grassland in the European Union in terms of utilisation, productivity, food security and the importance of adoption of technical innovations in increasing sustainability of pasture-based ruminant production systems Northwest Europe: current practices and main agronomic and environmental. Sustainable meat and milk production from grasslands Volume 23 Grassland Science in Europe.
- Dumont, B., Rossignol, N., Decuq, F., Note, P., & Farruggia, A. (2020). How does pasture size alter plant-herbivore interactions among grazing cattle?. *Grass and Forage Science*, 75(4), 438-446. Doi: 10.1111/gfs.12503.
- Elorriaga, Y., Marino, D. J., Carriquiriborde, P., & Alicia, R. (2013). Screening of pharmaceuticals in surface water bodies of the Pampas region of Argentina. *Int. J. Environ. Res. Public Health*, 6, 330-339. Doi: 10.1504/IJENVH.2013.056974.
- Enriquez-Hidalgo, D., Teixeira, D. L., Lewis, E., Buckley, F., Boyle, L., & O'Driscoll, K. (2018). Behavioural responses of pasture based dairy cows to short term management in tie-stalls. *Applied Animal Behaviour Science*, *198*, 19-26. Doi: 10.1016/j.applanim.2017.09.012.
- Fleurance, G., Farruggia, A., Lanore, L., & Dumont, B. (2016). How does stocking rate influence horse behaviour, performances and pasture biodiversity in mesophile grasslands?. *Agriculture, Ecosystems and Environment*, 231, 255-263. Doi: 10.1016/j.agee.2016.06.044.
- Fracchiolla, M., Terzi, M., D'Amico, F. S., Tedone, L., & Cazzato, E. (2017). Conservation and pastoral value of former arable lands in the agro-pastoral system of the Alta Murgia national park (Southern Italy). *Italian Journal of Agronomy*, *12*(2). Doi: 10.4081/ija.2017.847.
- Fraser, M. D., Vallin, H. E., & Roberts, B. P. (2022). Animal board invited review: Grassland-based livestock farming and biodiversity. *Animal*, 16(12). Doi: 10.1016/j.animal.2022.100671.
- Freschi, P., Musto, M., Paolino, R., & Cosentino, C. (2015). Grazing and biodiversity conservation: Highlights on a natura 2000 network site. In: *The Sustainability of Agro-Food and Natural Resource Systems in the Mediterranean Basin* (pp. 271-288). Springer International Publishing. Doi: 10.1007/978-3-319-16357-4_18.
- Futa, B., Ukalska-Jaruga, A., Tajchman, K., Janiszewski, P., & Pecio, M. (2024). Variability in Nutrient Content and Biochemical Parameters of Soil Under Rotational Pasture Management of Farmed Fallow Deer. *Agriculture (Switzerland)*, 14(11). Doi: 10.3390/agriculture14112011.
- Gallé, R., Urák, I., Nikolett, G. S., & Hartel, T. (2017). Sparse trees and shrubs confers a high biodiversity to pastures: Case study on spiders from Transylvania. *PLoS ONE*, *12*(9). Doi: 10.1371/journal.pone.0183465.

- Grodkowski, G., Gołębiewski, M., Slósarz, J., Grodkowska, K., Kostusiak, P., Sakowski, T., & Puppel, K. (2023). Organic Milk Production and Dairy Farming Constraints and Prospects under the Laws of the European Union. *Animals*, 13(9). MDPI. Doi: 10.3390/ani13091457.
- Hennessy, D., Delaby, L., van den Pol-van Dasselaar, A., & Shalloo, L. (2020). Increasing grazing in dairy cow milk production systems in Europe. *Sustainability (Switzerland)*, 12(6). Doi: 10.3390/su12062443.
- Hesselmann, M., Thorne, S., Vitra, A., Steiner, A. K., Leiber, F., & Dittmann, M. T. (2025). Foraging preferences of dairy cows grazing on contrasted multispecies swards. *Veterinary and Animal Science*, 28. Doi: 10.1016/j.vas.2025.100439.
- Hessle, A., & Danielsson, R. (2024). Cattle population required for favorable conservation status of management-dependent semi-natural grasslands and forests, and associated increase in enteric methane emissions. *Journal for Nature Conservation*, 78. Doi: 10.1016/j.jnc.2024.126571.
- Jechalke, S., Heuer, H., Siemens, J., Amelung, W., and Smalla, K. (2014). Fate and effects of veterinary antibiotics in soil. *Trends Microbiol.*, 22, 536-545. Doi: 10.1016/j.tim.2014.05.005.
- Kulik, M., Tajchman, K., Lipiec, A., Bąkowski, M., Ukalska-Jaruga, A., Ceacero, F., Pecio, M., & Steiner-Bogdaszewska, Ż. (2023). The Impact of Rotational Pasture Management for Farm-Bred Fallow Deer (Dama dama) on Fodder Quality in the Context of Animal Welfare. *Agronomy*, 13(4). Doi: 10.3390/agronomy13041155.
- Larraz, V., Barrantes, O., & Reiné, R. (2024). Habitat Selection by Free-Grazing Sheep in a Mountain Pasture. *Animals*, *14*(13). Doi: 10.3390/ani14131871.
- Lennartsson, T., & Westin, A. (2024). Can Current Grazing Practices Preserve Biodiversity in Semi-natural Pastures? A Study of the Historical Ecology of Swedish Infield Pastures. *Martor*, 29, 46-67. Doi: 10.57225/martor.2024.29.04.
- Marchegiani, S., Gislon, G., Marino, R., Caroprese, M., Albenzio, M., Pinchak, W. E., Carstens, G. E., Ledda, L., Trombetta, M. F., Sandrucci, A., Pasquini, M., Deligios, P. A., & Ceccobelli, S. (2025). Smart technologies for sustainable pasture-based ruminant systems: A review. Smart Agricultural Technology, 10. Doi: 10.1016/j.atech.2025.100789.
- McCormick, I. A., & Stokes, J. E. (2023). Stakeholder Challenges and Opportunities of GPS Shock Collars to Achieve Optimum Welfare in a Conservation or Farm Setting. *Animals*, *13*(19). Doi: 10.3390/ani13193084.
- Minea, G., Mititelu-Ionuş, O., Gyasi-Agyei, Y., Ciobotaru, N., & Rodrigo-Comino, J. (2022). Impacts of Grazing by Small Ruminants on Hillslope Hydrological Processes: A Review of European Current Understanding. *Water Resources Research* 2024 58(3). Doi: 10.1029/2021WR030716.
- Molle, G., Cannas, A., & Gregorini, P. (2022). A review on the effects of part-time grazing herbaceous pastures on feeding behaviour and intake of cattle, sheep and horses. *Livestock Science*2024 263. Doi: 10.1016/j.livsci.2022.104982.
- Morais, T. G., dos Santos, M. P., Barão, L., Domingos, T., & Teixeira, R. F. M. (2025). Grazing or confining Decoding Beef's environmental footprint. Environmental Impact Assessment Review, 112. Doi: 10.1016/j.eiar.2025.107846.

- Morais, T. G., Teixeira, R. F. M., Rodrigues, N. R., & Domingos, T. (2018). Carbon footprint of milk from pasture-based dairy farms in Azores, Portugal. *Sustainability (Switzerland)*, 10(10). Doi: 10.3390/su10103658.
- Moret-Fernández, D., Arroyo, A. I., Herrero, J., Barrantes, O., Alados, C. L., & Pueyo, Y. (2021). Livestock grazing effect on the hydraulic properties of gypseous soils in a Mediterranean region. *Catena*, 207. Doi: 10.1016/j.catena.2021.105697.
- Moret-Fernández, D., Pueyo, Y., Bueno, C. G., & Alados, C. L. (2011). Hydrophysical responses of gypseous and non-gypseous soils to livestock grazing in a semi-arid region of NE Spain. *Agricultural Water Management*, *98*(12), 1822-1827. Doi: 10.1016/j.agwat.2011.07.001.
- Niero, G., Bobbo, T., Callegaro, S., Visentin, G., Pornaro, C., Penasa, M., Cozzi, G., de Marchi, M., & Cassandro, M. (2021). Dairy cows' health during alpine summer grazing as assessed by milk traits, including differential somatic cell count: A case study from Italy. *Animals*, 11(4). Doi: 10.3390/ani11040981.
- O'Brien, D., Bohan, A., McHugh, N., & Shalloo, L. (2016). A life cycle assessment of the effect of intensification on the environmental impacts and resource use of grass- based sheep farming. *Agricultural Systems*, *148*, 95-104. Doi: 10.1016/j. agsy.2016.07.004.
- O'Driscoll, K., Lewis, E., & Kennedy, E. (2019). Effect of feed allowance at pasture on the lying behaviour of dairy cows. *Applied Animal Behaviour Science*, *213*, 40-46. Doi: 10.1016/j.applanim.2019.02.002.
- O'Leary, N. W., Byrne, D. T., Garcia, P., Werner, J., Cabedoche, M., & Shalloo, L. (2020). Grazing cow behavior's association with mild and moderate lameness. *Animals*, 10(4). Doi: 10.3390/ani10040661.
- Opio, C., Gerber, P., Mottet, A., Falculli, A., Tempio, G., MacLeod, M. et al. (2013). Greenhouse gas emissions from ruminant supply chains - A global life cycle assessment. Food and Agriculture Organization of the United Nations (FAO), 214.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372(71). Doi: 10.1136/bmj.n71.
- Pauler, C. M., Homburger, H., Lüscher, A., Scherer-Lorenzen, M., & Schneider, M. K. (2025). Ecosystem services in mountain pastures: A complex network of site conditions, climate and management. Agriculture, Ecosystems and Environment, 377. Doi. 10.1016/j.agee.2024.109272.
- Phelps, L. N., & Kaplan, J. O. (2017). Land use for animal production in global change studies: Defining and characterizing a framework. *Global Change Biology*, 23(11), 4457-4471. Doi: 10.1111/gcb.13732.
- Plieninger, T., Hartel, T., Martín-López, B., Beaufoy, G., Bergmeier, E., Kirby, K., Montero, M. J., Moreno, G., Oteros-Rozas, E., & van Uytvanck, J. (2015). Woodpastures of Europe: Geographic coverage, social-ecological values, conservation management, and policy implications. *Biological Conservation*, 190, 70-79. Doi: 10.1016/j.biocon.2015.05.014.
- Ponzetta, M. P., Grohmann, F., & Argenti, G. (2010). Effects of two restoration practices on the vegetation of a semi-natural grassland invaded by bracken in Central Italy. *Italian Journal of Agronomy*, 5(3), 233-240. Doi: 10.4081/ija.2010.233.

- Potenza, C., & Fedele, V. (2011). Effectiveness of the GAEC cross-compliance standard protection of permanent pasture in relation to grazing and pasture conservation management in marginal mountain areas. *Italian Journal of Agronomy*, 6(Supl.1), 83-86. Doi: 10.4081/ija.2011.6.s1.e11.
- Ripamonti, A., Mantino, A., Annecchini, F., Cappucci, A., Casarosa, L., Turini, L., Foggi, G., & Mele, M. (2023). Outcomes of a comparison between pastoral and silvopastoral management on beef cattle productivity, animal welfare and pasture depletion in a Mediterranean extensive farm. *Agroforestry Systems*, *97*(6), 1071-1086. Doi: 10.1007/s10457-023-00848-w.
- Ripoll-Bosch, R., de Boer, I. J. M., Bernués, A., & Vellinga, T. v. (2013). Accounting for multi-functionality of sheep farming in the carbon footprint of lamb: A comparison of three contrasting Mediterranean systems. *Agricultural Systems*, 116, 60-68. Doi: 10.1016/j.agsy.2012.11.002.
- Rivera, D., Verde, A., Fajardo Rodríguez, J., Ríos, S., Alcaraz, F., Cárceles, C., Ortíz, J., Valdés, A., Ruíz-Gallardo, J. R., García-Flores, A., Palazón, J. A., & Obón, C. (2022). Ethnoveterinary Medicine and Ethnopharmacology in the Main Transhumance Areas of Castilla-La Mancha (Spain). *Frontiers in Veterinary Science*, 9. Doi: 10.3389/fvets.2022.866132.
- Rodriguez, J. (2024). One Health Ethics and the Ethics of Zoonoses: A Silent Call for Global Action. *Veterinary Sciences*, 11(9), 394. Doi: 10.3390/vetsci11090394.
- Roeder, N., Lederbogen, D., Trautner, J., Bergamini, A., Stofer, S., & Scheidegger, C. (2010). The impact of changing agricultural policies on jointly used rough pastures in the Bavarian Pre-Alps: An economic and ecological scenario approach. *Ecological Economics*, 69(12), 2435-2447. Doi: 10.1016/j. ecolecon.2010.07.013.
- Rutherford, N. H., Lively, F. O., & Arnott, G. (2021). A Review of Beef Production Systems for the Sustainable Use of Surplus Male Dairy-Origin Calves Within the UK. *Frontiers in Veterinary Science*, 8. Doi: 10.3389/fvets.2021.635497.
- Sala, V., Moja, L., Moschetti, I., Bidoli, S., Pistotti, V., & Liberati, A. (n.d.). Revisioni sistematiche-Breve guida all'uso.
- Sargeant, J. M., Amezcua, M. D. R., Rajic, A., & Waddell, L. (2005). *Guide to conducting systematic reviews in agri-food public health*. Food Safety Research and Response Network. Supported by the United States Department of Agriculture, Cooperative Research, Education and Extension Service.
- Schmitz, A., & Isselstein, J. (2020). Effect of grazing system on grassland plant species richness and vegetation characteristics: Comparing horse and cattle grazing. *Sustainability (Switzerland)*, *12*(8). Doi: 10.3390/SU12083300.
- Simionesei, L., Ramos, T. B., Oliveira, A. R., Jongen, M., Darouich, H., Weber, K., Proença, V., Domingos, T., & Neves, R. (2018). Modeling soilwater dynamics and pasture growth in the montado ecosystem using MOHID land. *Water (Switzerland)*, 10(4). Doi: 10.3390/w10040489.
- Sitzia, T., Semenzato, P., & Trentanovi, G. (2010). Natural reforestation is changing spatial patterns of rural mountain and hill landscapes: A global overview. In *Forest Ecology and Management*, 259(8), 1354-1362). Doi: 10.1016/j. foreco.2010.01.048.

- Smetanová, A., Follain, S., David, M., Ciampalini, R., Raclot, D., Crabit, A., & le Bissonnais, Y. (2019). Landscaping compromises for land degradation neutrality: The case of soil erosion in a Mediterranean agricultural landscape. *Journal of Environmental Management*, 235, 282-292. Doi: 10.1016/j.jenvman.2019.01.063.
- Steiner, M., Öckinger, E., Karrer, G., Winsa, M., & Jonsell, M. (2016). Restoration of semi-natural grasslands, a success for phytophagous beetles (Curculionidae). *Biodiversity and Conservation*, 25(14), 3005-3022. Doi: 10.1007/s10531-016-1217-4.
- Strand, G. H., Hansen, I., de Boon, A., & Sandström, C. (2019). Carnivore Management Zones and their Impact on Sheep Farming in Norway. *Environmental Management*, 64(5), 537-552. Doi: 10.1007/s00267-019-01212-4.
- Szyszko-Podgórska, K., Szweda, Ż., Świątek, M., Ukalska, J., Pietrasz, K., Pietrasz, M., Wilk, P., Orlińska-Woźniak, P., Szalińska, E., Rokicki, T., Tylkowski, S., & Niżnikowski, R. (2024). Impact of Land Use on Peat Soil Elemental Content and Carabidae and Plant Species Composition and Abundance. *Sustainability (Switzerland)*, *16*(11). Doi: 10.3390/su16114420.
- Tittonell, P. (2021). Beyond CO2: Multiple ecosystem services from ecologically intensive grazing landscapes of South America. *Frontiers in Sustainable Food Systems*, 5. Doi: 10.3389/fsufs.2021.664103.
- Tölgyesi, C., Kelemen, A., Bátori, Z., Kiss, R., Hábenczyus, A. A., Havadtői, K., Varga, A., Erdős, L., Frei, K., Tóth, B., & Török, P. (2023). Maintaining scattered trees to boost carbon stock in temperate pastures does not compromise overall pasture quality for the livestock. *Agriculture, Ecosystems and Environment*, 351. Doi: 10.1016/j.agee.2023.108477.
- Tonn, B., Densing, E. M., Gabler, J., & Isselstein, J. (2019). Grazing-induced patchiness, not grazing intensity, drives plant diversity in European low-input pastures. *Journal of Applied Ecology*, *56*(7), 1624-1636. Doi: 10.1111/1365-2664.13416.
- Toro-Mujica, P. M., Aguilar, C., Vera, R., Barba, C., Rivas, J., & García-Martínez, A. (2015). Changes in the pastoral sheep systems of semi-arid Mediterranean areas: Association with common agricultural policy reform and implications for sustainability. *Spanish Journal of Agricultural Research*, *13*(2), 1-11. Doi: 10.5424/sjar/2015132-6984.
- Tsiobani, E. T., Yiakoulaki, M. D., Hasanagas, N. D., & Antoniou, I. E. (2023). Water buffalo social networks during their activities on pasture under the rain. *Turkish Journal of Veterinary and Animal Sciences*, 47(6), 576-583. Doi: 10.55730/1300-0128.4327.
- Turini, L., Foggi, G., Mantino, A., Gasparoni, E., Vichi, F., Silvi, A., Armenia, G., Sala, G., Bonelli, F., Sgorbini, M., & Mele, M. (2025). Changes in hematological and hematochemical parameters in lactating dairy sheep according to different pasture management and heat stress risk: A longitudinal study. *Veterinary and Animal Science*, 27. Doi: 10.1016/j.vas.2024.100419.
- van Klink, R., Nolte, S., Mandema, F. S., Lagendijk, D. D. G., WallisDeVries, M. F., Bakker, J. P., Esselink, P., & Smit, C. (2016). Effects of grazing management on biodiversity across trophic levels The importance of livestock species and stocking density in salt marshes. *Agriculture, Ecosystems and Environment*, 235, 329-339. Doi: 10.1016/j.agee.2016.11.001.

- Vargas-Canales, J. M., Orozco-Cirilo, S., Estrada, S., del Carpio-Ovando, P. S., Camacho-Vera, J. H., López-Carmona, D., García-Melchor, N., Rodríguez-Haros, B., Valdés-Cobos, A., Sánchez-Torres, Y., Fresnedo-Ramírez, J., Palacios-Rangel, M. I., Ocampo-Ledesma, J. G., Barrera-Perales, O. T., Pineda-Pineda, J., Kreimer, P., García-Cruz, J. C., Reyes-Barrera, D. M., Montiel-Flores, J. C., ... de Gortari-Rabiela, R. (2024). Science, technology, agri-food systems, health, and wellbeing: logic, dynamics, and relationships. Frontiers in Sustainable Food Systems, 8. Doi: 10.3389/fsufs.2024.1344357.
- Verduna, T., Blanc, S., Merlino, V. M., Cornale, P., & Battaglini, L. M. (2020). Sustainability of Four Dairy Farming Scenarios in an Alpine Environment: The Case Study of Toma di Lanzo Cheese. *Frontiers in Veterinary Science*, 7. Doi: 10.3389/fyets.2020.569167.
- Wróbel, B., Zielewicz, W., & Staniak, M. (2023). Challenges of Pasture Feeding Systems Opportunities and Constraints. *Agriculture (Switzerland)*, 13(5). Doi: 10.3390/agriculture13050974.
- Zhu, S., Olde, L., Lewis, K., Quaife, T., Cardenas, L., Loick, N., Xu, J., & Hill, T. (2023). Eddy covariance fluxes over managed ecosystems extrapolated to field scales at fine spatial resolutions. *Agricultural and Forest Meteorology*, 342. Doi: 10.1016/j.agrformet.2023.109675.

Giulia Pastorelli

Researcher, CREA Research Centre for Agricultural Policies and Bioeconomy, Rome (Italy)

E-mail: giulia.pastorelli@crea.gov.it

Based at the MASAF technical office within the National Rural Network, she contributes to rural development policies relating to the implementation of the Common Agricultural Policy (CAP) in Italy. Her research focuses on the economic analysis of CAP measures and specifically on animal welfare, organic farming, the circular economy, and biodiversity conservation.

https://orcid.org/0009-0002-3295-0302

Ilaria Falconi

Research technologist, CREA Research Centre for Agricultural Policies and Bioeconomy, Rome (Italy)

E-mail: ilaria.falconi@crea.gov.it

She is a member of the expert group of the European carbon farming. She is a member of the ENRD group of experts on carbon farming. She is a member of the working groups for the National Ecological Transition Plan. She is a member of the EU expert group "Sustainability and quality in agriculture and rural areas - Methane emissions subgroup".

Maria Assunta D'Oronzio

Senior researcher, CREA Research Centre for Agricultural Policies and Bioeconomy, Potenza (Italy)

E-mail: massunta.doronzio@crea.gov.it

She contributes to rural development policies related to the implementation of the Common Agricultural Policy (CAP) in Italy. Her research focuses on the economic analysis of supply chains, the sustainability of agricultural production, precision and digital agriculture, the circular economy and the conservation of biodiversity. She is the author and co-author of numerous publications and has presented at various national and international conferences.

Raffaella Pergamo

Senior researcher, CREA Research Centre for Agricultural Policies and Bioeconomy, Rome (Italy)

E-mail: raffaella.pergamo@crea.gov.it

Based at MASAF's technical office within the National Rural Network, she contributes to rural development policies related to the implementation of the Common Agricultural Policy (CAP) in Italy. Her research focuses on the economic analysis of supply chains, the sustainability of agricultural production, precision and digital agriculture, the circular economy and the conservation of biodiversity. She is the author and co-author of numerous publications and has presented at various national and international conferences.