
The microbiota's funnel effect: How multiple factors independently converge in changing the microbiota

Massimo Agnoletti*, Alessio Fasano**

Received 13 February, revised 15 February, accepted 18 February

Online first: 25 February 2025

* University of Verona.

info@massimoagnoletti.it

** Harvard Medical School.

afasano@mgh.harvard.edu

Please cite: Agnoletti M., Fasano A. (2025). The microbiota's funnel effect: How multiple factors independently converge in changing the microbiota. *PNEI Review*. DOI: 10.3280/pnei2025oa19441

Abstract: *The science of the microbiota, by revealing the complexity of interactions between the world of microorganisms that colonize us and that of human cells, represents the dawn of a new paradigm in biomedical and psychological sciences. It allows for a more complex yet more promising perspective on human health and disease compared to previous approaches. Scientific literature has now identified specific lifestyle-related factors (nutrition, environmental physical-chemical quality, psychological well-being, etc.) that significantly influence the composition of the microbiota. This paper introduces the concept of the “funnel effect” of the microbiota to describe the convergent and partially independent nature of these factors (nutrition, physical activity, psychological well-being, sleep quality, social support, environmental physical-chemical quality, circadian rhythms) on the composition of the intestinal microbiota, thereby impacting the overall health of the human organism. The “funnel effect” of the microbiota has some highly relevant clinical implications, emphasizing the need for an integrated psycho-neuro-endocrine-immunological approach, in contrast to the highly specialized and molecularly focused approach that is currently widely adopted.*

Key words: *Microbiota, PNEI paradigm, Nutrition, Physical activity, Psychological well-being, Epigenetic.*

Introduction

At the end of the last century, the massive Human Genome Project aimed to fully map human DNA with the ambition of understanding and potentially solving any human disease. This optimistic expectation stemmed from the theoretical concept of the so-called “central dogma of molecular biology”, which posits that a gene (part of the genotype) corresponds to a specific protein (phenotype) and that there can be no informational flow from the phenotype to the genotype – only a unidirectional flow from the gene to the protein structure (Bottaccioli F. & Bottaccioli A.G., 2017; Gottlieb, 2000). Paradoxically, the success of the Human Genome Project in fully sequencing human DNA marked the end of the “central dogma of molecular biology” because it exposed the theoretical inadequacy of this concept considering the data emerging from genetic analysis and the complexity of human bio-psycho-social interactions. A striking example is the fact that a complex organism like a human possesses a genetic heritage of “only” about 25,000 genes, compared to wheat, which has around 150,000, and that the percentage of coding genes is limited to approximately 2% of the entire human genome.

As often happens in the history of science, the partial failure of the ambitious Human Genome Project, on one hand, led to an increasing awareness of the explanatory inadequacy of the “central dogma of molecular biology” in relation to human phenotypic complexity. On the other hand, however, it laid the foundation for two major revolutions – still ongoing – in the scientific landscape. The first revolution concerns epigenetics, the study of factors (both hereditary and non-hereditary) that modify an organism’s phenotype without altering the DNA sequence. In other words, it involves studying the flow of information from the environment to the DNA, thereby influencing the expression of the genotype. The second revolution is represented by microbiota science, which examines the trillions of microorganisms (bacteria, fungi, viruses, protozoa) that colonize our bodies.

The effects of microbiota

Thanks to the genetic analysis technologies developed for the Human Genome Project, significant progress has been made in studying the microbiota and understanding the extent to which interactions among its microorganisms affect human cells. This emerging field of study has begun to uncover the astonishing complexity and significant impact of the microorganisms inhabiting our bodies in shaping human health and quality of life. Strengthening this broader perspective – one that includes the microbiota’s influence on human bio-psycho-social well-being – there is now substantial evidence showing that the microbial ecosystem colonizing us pro-

foundly affects complex functions such as nutrient absorption, immune system regulation, and neurodevelopment. It modulates systems like the stress axis and even influences cognitive, emotional, and motivational aspects. The interaction between the microbiota and intestinal permeability plays a particularly crucial role in modulating the immune system and the epigenetic impact on the human host. This makes it a key factor in the development of chronic inflammatory diseases, which have become a true “epidemic” in industrialized countries (Fasano, 2020).

The remarkable influence of the microbiota has shed light on the etiopathogenesis of conditions such as celiac disease, obesity, and ulcerative colitis, as well as psychological disorders, including anxiety, depression, and various psychopathologies such as autism spectrum disorders and schizophrenia (Caio *et al.*, 2019; Cheung *et al.*, 2019; Foster & McVey Neufeld, 2013; Garrett *et al.*, 2007; Li & Zhou, 2016; Mangiola *et al.*, 2016; Rodrigues-Amorim *et al.*, 2018; Sharon *et al.*, 2019; Simpson *et al.*, 2021).

It is important to note that the emergence of the new paradigm, which positions the microbiota as a key player in our health, has been driven precisely by research demonstrating how the microbiota influences neurobehavioral and psychological dimensions – such as anxious or depressive states, sociality, and risk perception (Allen *et al.*, 2017; Ann *et al.*, 2024; Bercik *et al.*, 2011; Bravo *et al.*, 2011; Carloni *et al.*, 2021; Chen *et al.*, 2019; Cheung *et al.*, 2019; Cryan & Dinan, 2012; Cryan & OMahony, 2011; De Palma *et al.*, 2015; Farmer, Randall & Aziz, 2014; Koenig *et al.*, 2011; Ottman *et al.*, 2012; Wu *et al.*, 2021) and how certain phenotypic characteristics of these dimensions can even be transferred from one organism to another through the so-called «microbiota transplantation» (Chinna Meyyappan *et al.*, 2020; Collins *et al.*, 2013; Cryan & Dinan, 2012; Kelly *et al.*, 2016).

It has long been known that the mind can influence intestinal well-being (also because this is intuitively evident from an experiential perspective), but the study of the microbiota has also demonstrated the existence of the opposite causal direction, in which intestinal microorganisms exert an effect on psychosocial dynamics.

The significant impact of the microbiota on the human organism includes endocrine and immunological dynamics, as well as neural and psychological ones, thus affecting the entire complexity of human bio-psycho-social interactions. This is one of the reasons why it would be more appropriate to refer to the “microbiota-gut-brain-mind axis” rather than the reductive, yet widely used, term “microbiota-gut-brain axis”, which erroneously considers the mind and brain to be entirely coincident (Agnoletti, 2023a).

The impact of microbiota interaction on human cells is remarkable, considering that the total mass of bacteria alone is estimated to be between 0.2 kg and 1 kg (70% of which are located in the gut), the number of bacteria is estimated to be equal to or greater than that of human cells, and most importantly, the genetic content of bacte-

ria alone is approximately 100 to 1.000 times greater than that of humans (Sender, Fuchs & Milo, 2016a; Sender, Fuchs & Milo, 2016b).

Given the symbiotic relationship between microbiota and human cells, the microbiome – the collective genetic heritage of the microorganisms that make up the microbiota – represents a variable component of the human genome. In fact, due to its significantly larger genetic contribution compared to the human genome, the microbiota plays a crucial epigenetic role in relation to human cells, making it fundamental in determining human well-being, health, and longevity (Chang *et al.*, 2014; Claesson *et al.*, 2012; Cornuti *et al.*, 2013; Dalile *et al.*, 2019; Kumar *et al.*, 2014; López-Otín *et al.*, 2013; Ottaviani, 2011).

Understanding epigenetic dynamics is essential to grasp the complex interaction between the microbiota and human cells because studying the factors that influence genetic expression allows us to conceptualize the entire microbiota ecosystem as an extra-genetic factor (“extra” in relation to human DNA) that further extends the adaptive capacity of human cells. The additional genetic contribution of the microbiota enables the human holobiont organism (represented by the collaboration between human cells and microorganisms) to adapt more effectively to different environmental conditions, providing clear evolutionary advantages (Agnoletti, 2023b; Gasbarrini, Dionisi & Gasbarrini, 2019; Fasano, 2022). The nature of the genetic contribution of the microbiota represents the greatest factor of interindividual diversity. In fact, while human interindividual genetic heritage is approximately 99,9% identical, the genetic diversity of the microbiota between two individuals can reach as high as 80-90% (Gasbarrini, Dionisi & Gasbarrini, 2019).

The genetic diversity of the microbiota is primarily determined by environmental factors and only marginally by the host’s human genetics (Rothschild *et al.*, 2018). Therefore, we can assert that almost the entire contribution of the microbiota to human cells is due to environmental factors rather than human genetic content. Human fitness is thus highly dependent on the epigenetic role of the microbiota, as it significantly expands the body’s ability to adapt to variable contexts, such as the food we eat, the physical activity we engage in, the physicochemical quality of the environments we frequent, and the psychosocial stress we experience. Despite these variations, the microbiota still ensures the homeostasis of key physiological systems, including blood pressure, heart rate, blood volume, hydration, pH, bone density, and more.

For the purposes of this discussion, it is important to note that the currently available literature has clearly demonstrated that factors such as:

- Nutrition (Putignani & Dallapiccola, 2016; Valitutti, Cucchiara & Fasano, 2019; Zhang & Zuo, 2018),
- Physical activity (Allen *et al.*, 2018; Aragón-Vela *et al.*, 2021; Mohr *et al.*, 2020),
- Sleep quality (Neroni *et al.*, 2021; Sen *et al.*, 2021; Smith *et al.*, 2019),

- Circadian rhythms (Birmingham *et al.*, 2023; Thaïss *et al.*, 2014; Voigt *et al.*, 2016),
- Physicochemical environmental quality (De Filippis *et al.*, 2024; Estevinho *et al.*, 2024; Mousavi *et al.*, 2022),
- Perceived social support (Cryan *et al.*, 2019; Kim *et al.*, 2021; Winter *et al.*, 2018),
- Psychological well-being (Chang *et al.*, 2024; Ge *et al.*, 2022; Ilchmann-Diounou & Ménard, 2020), influence the composition of the microbiota, thereby modifying its adaptive contribution – both epigenetic and non-epigenetic – to the entire organism, ultimately affecting human health and quality of life. For the purposes of this discussion, it is particularly interesting to note that the currently available scientific literature does not highlight the dominance of one factor over the others in influencing the microbiota. Instead, it describes a dose-dependent quantitative effect for each of them.

All the factors considered influence one another to some extent (for instance, physical activity affects sleep quality, and circadian rhythms impact the metabolic effects of nutrition). However, each factor also maintains a certain degree of autonomy from the others due to its unique causal mechanism in shaping the microbiota. For example, the management of psychological stress and the physicochemical environmental quality are both factors that influence the microbiota, but they do so at least partially independently of each other, as they follow different causal pathways. The “sensitivity” of the microbiota to multiple factors is thus somewhat independent of the specific nature of the influencing factor itself. In other words, whether it is nutrition, psychological well-being, or physical activity, there will always be an impact on the composition of the microbiota. For instance, dysbiosis may initially stem from a nutritional deficiency, a psychological trauma, or excessive sedentary behavior, even though the causal dynamics linking nutrition, psychological states, and physical activity to the microbiota are at least partially independent of one another.

The fact that various factors converge in modifying the microbiota does not mean that their impact is the same in terms of intensity or frequency. Naturally, the causal pathway resulting from chronic sleep deprivation has a different impact on the microbiota compared to, for example, a single mild exposure to pesticides.


The funnel effect

Considering the diversity and partial independence of the different factors that influence the microbiota, it is equally interesting to highlight the common convergence in modifying the composition of this complex ecosystem that colonizes our body.

For these reasons, the authors propose the concept of the “microbiota funnel effect” to describe the characteristic convergence of the seven main factors identified in the scientific literature (nutrition, physical activity, psychological, sleep-related, so-

cial, environmental, and circadian), which, in a partially independent manner, contribute to altering the composition of the gut microbiota (see Figure 1).

Fig. 1. The Microbiota's Funnel Effect



The common convergence of the seven lifestyle-related factors on the microbiota has significant implications for peoples well-being and health, highlighting the need for a systemic and holistic approach to promoting well-being and treating suboptimal and pathological conditions.

From a clinical perspective, the “microbiota funnel effect” asserts that, given the high psycho-neuro-endocrine-immunological integration of the considered factors and the significant epigenetic impact of the microbiota on human cell function, the only way to ensure an effective clinical intervention is to adopt a systemic vision that includes a comprehensive evaluation of all factors. Continuing to use a specialist approach that analyses and treats only one (or even some, but not all) of these factors by decontextualizing it from the others is equivalent to adopting a reductionist perspective. This reductionistic approach, by failing to account for the convergent and partially independent nature described by the “microbiota funnel effect”, proves to be clinically ineffective, particularly in addressing chronic conditions.

The concept of the “microbiota funnel effect” calls for a radical shift in clinical protocols, as it necessitates at least an initial comprehensive (but not generic) as-

assessment of the individual factors involved and the patient's gut health (which includes the general state of the microbiota). For any professional whose goal is to enhance human health and well-being, the “microbiota funnel effect” represents a new paradigm that demonstrates the clinical necessity of a truly integrated, holistic, and scientific approach – essential for addressing complex issues involving the microbiota.

Author Contributions: Conceptualization: M.A., A.F.; writing – original draft preparation: M.A.; writing – review M.A., A.F.; editing: M.A.

Funding: This research received no external funding. Conflicts of Interest: The authors declare no conflict of interest. All authors have read and agreed to the published version of the manuscript.

References

- Agnoletti M. (2023a).** Il concetto di Adattoma nel paradigma di benessere e salute Bio-Psico-Sociale. *Medicalive Magazine*, 1, 16-20.
- Agnoletti M. (2023b).** Perché si dovrebbe parlare di asse microbiota-intestino-cervello-mente. *Medicalive Magazine*, 6, 12-16.
- Allen A.P., Dinan T. G., Clarke G., & Cryan J.F. (2017).** A psychology of the human brain-gut-microbiome axis. *Social and Personality Psychology Compass*, 11(4), e12309. <https://doi.org/10.1111/spc3.12309>
- Allen J.M., Mailing L.J., Niemi G.M., Moore R., Cook M.D., White B.A., Holscher H.D., & Woods J.A. (2018).** Exercise alters gut microbiota composition and function in lean and obese humans. *Medicine and Science in Sports and Exercise*, 50(4), 747–757. <https://doi.org/10.1249/MSS.0000000000001495>
- An E., Delgado D.R., & Yang J. et al. (2024).** Stress-resilience impacts psychological wellbeing as evidenced by brain–gut microbiome interactions. *Nature Mental Health*, 2, 935–950. <https://doi.org/10.1038/s44220-024-00266-6>
- Aragón-Vela J., Solis-Urra P., Ruiz-Ojeda F.J., Álvarez-Mercado A.I., Olivares-Arancibia J., & Plaza-Diaz J. (2021).** Impact of exercise on gut microbiota in obesity. *Nutrients*, 13(11), 3999. <https://doi.org/10.3390/nu13113999>
- Bercik P., Denou E., Collins J., Jackson W., Lu J., Jury J., Deng Y., Blennerhassett P., Macri J., McCoy K.D., Verdu E.F., & Collins S.M. (2011).** The intestinal microbiota affects central levels of brain-derived neurotropic factor and behavior in mice. *Gastroenterology*, 141(2), 599–609.e6093. <https://doi.org/10.1053/j.gastro.2011.04.052>
- Bermingham K.M., Stensrud S., Asnicar F., Valdes A.M., Franks P.W., Wolf J., Hadjigeorgiou G., Davies R., Spector T.D., Segata N., Berry S.E., & Hall W.L. (2023).** Exploring the relationship between social jetlag with gut microbial composition, diet and cardiometabolic health, in the ZOE PREDICT 1 cohort. *European Journal of Nutrition*, 62(8), 3135–3147. <https://doi.org/10.1007/s00394-023-03204-x>
- Bottaccioli F. & Bottaccioli A.G. (2017).** *Psiconeuroendocrinologia e scienza della cura integrata. Il Manuale*. EDRA.
- Bravo J.A., Forsythe P., Chew M.V., Escaravage E., Savignac H.M., Dinan T.G., Bienenstock J., & Cryan J.F. (2011).** Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proceedings of the National Academy of Sciences of the United States of America*, 108(38), 16050–16055. <https://doi.org/10.1073/pnas.1102999108>
- Caio G., Volta U., Sapone A., Leffler D.A., De Giorgio R., Catassi C., & Fasano A. (2019).** Celiac disease: A comprehensive current review. *BMC Medicine*, 17(1), 142. <https://doi.org/10.1186/s12916-019-1380-z>
- Carloni S., Bertocchi A., Mancinelli S., Bellini M., Erreni M., Borreca A., Braga D., Giugliano S., Mozzarelli A.M., Manganaro D., Fernandez Perez D., Colombo F., Di Sabatino A., Pasini D., Penna G., Matteoli**

- M., Lodato S., & Rescigno M. (2021). Identification of a choroid plexus vascular barrier closing during intestinal inflammation. *Science*, 374(6566), 439–448. <https://doi.org/10.1126/science.abc6108>
- Chang H., Perkins M.H., Novaes L.S., Qian F., Zhang T., Neckel P.H., Scherer S., Ley R.E., Han W., & de Araujo I.E. (2024). Stress-sensitive neural circuits change the gut microbiome via duodenal glands. *Cell*, 187(19), 5393–5412.e30. <https://doi.org/10.1016/j.cell.2024.07.019>
- Chang P.V., Hao L., Offermanns S., & Medzhitov R. (2014). The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proceedings of the National Academy of Sciences of the United States of America*, 111(6), 2247–2252. <https://doi.org/10.1073/pnas.1322269111>
- Chen Y.H., Bai J., Wu D., Yu S.F., Qiang X. L., Bai H., Wang H.N., & Peng Z.W. (2019). Association between fecal microbiota and generalized anxiety disorder: Severity and early treatment response. *Journal of Affective Disorders*, 259, 56–66. <https://doi.org/10.1016/j.jad.2019.08.014>
- Cheung S.G., Goldenthal A.R., Uhlemann A.C., Mann J.J., Miller J.M., & Sublette M.E. (2019). Systematic review of gut microbiota and major depression. *Frontiers in Psychiatry*, 10, 34. <https://doi.org/10.3389/fpsy.2019.00034>
- Chinna Meyyappan A., Forth E., Wallace C. J. K., & Milev R. (2020). Effect of fecal microbiota transplant on symptoms of psychiatric disorders: A systematic review. *BMC Psychiatry*, 20(1), 299. <https://doi.org/10.1186/s12888-020-02654-5>
- Claesson M.J., Jeffery I.B., Conde S., Power S.E., O'Connor E.M., Cusack S., Harris H.M., Coakley M., Lakshminarayanan B., O'Sullivan O., et al. (2012). Gut microbiota composition correlates with diet and health in the elderly. *Nature*, 488, 178–184. <https://doi.org/10.1038/nature11319>
- Collins S., Kassam Z., & Bercik P. (2013). The adoptive transfer of behavioral phenotype via the intestinal microbiota: Experimental evidence and clinical implications. *Current Opinion in Microbiology*, 16, 240–245. <https://doi.org/10.1016/j.mib.2013.06.004>
- Cornuti S., Chen S., Lupori L., Finamore F., Carli F., Samad M., Fenizia S., Caldarelli M., Damiani F., Raimondi F., Mazziotti R., Magnan C., Rocchiccioli S., Gastaldelli A., Baldi P., & Tognini P. (2023). Brain histone beta-hydroxybutyrylation couples metabolism with gene expression. *Cellular and Molecular Life Sciences*, 80(1), 28. <https://doi.org/10.1007/s00018-022-04673-9>
- Cryan J.F., O'Riordan K. J., Cowan C., Sandhu K.V., Bastiaansen T., Boehme M., Codagnone M.G., Cusotto S., Fulling C., Golubeva A.V., Guzzetta K.E., Jaggar M., Long-Smith C.M., Lyte J.M., Martin J.A., Molinero-Perez A., Moloney G., Morelli E., Morillas E., O'Connor R., & Dinan T.G. (2019). The microbiota-gut-brain axis. *Physiological Reviews*, 99(4), 1877–2013. <https://doi.org/10.1152/physrev.00018.2018>
- Cryan J.F., & Dinan T. G. (2012). Mind-altering microorganisms: The impact of the gut microbiota on brain and behaviour. *Nature Reviews Neuroscience*, 13(10), 701–712. <https://doi.org/10.1038/nrn3346>
- Cryan J.F., & O'Mahony S.M. (2011). The microbiome-gut-brain axis: From bowel to behavior. *Neurogastroenterology and Motility*, 23(3), 187–192. <https://doi.org/10.1111/j.1365-2982.2010.01664.x>
- Dalile B., Van Oudenhove L., Vervliet B., & Verbeke K. (2019). The role of short-chain fatty acids in microbiota-gut-brain communication. *Nature Reviews Gastroenterology & Hepatology*, 16(8), 461–478. <https://doi.org/10.1038/s41575-019-0157-3>
- De Palma G., Blennerhassett P., Lu J., Deng Y., Park A.J., Green W., Denou E., Silva M.A., Santacruz A., Sanz Y., Surette M.G., Verdu E.F., Collins S.M., & Bercik P. (2015). Microbiota and host determinants of behavioural phenotype in maternally separated mice. *Nature Communications*, 6, 7735. <https://doi.org/10.1038/ncomms8735>
- Estevinho M.M., Midya V., Cohen-Mekelburg S., Allin K.H., Fumery M., Pinho S.S., Colombel J.F., & Agrawal M. (2024). Emerging role of environmental pollutants in inflammatory bowel disease risk, outcomes and underlying mechanisms. *Gut*. Advance online publication. <https://doi.org/10.1136/gutjnl-2024-332523>
- Farmer A.D., Randall H.A., & Aziz Q. (2014). It's a gut feeling: how the gut microbiota affects the state of mind. *The Journal of Physiology*, 592(14), 2981–2988. <https://doi.org/10.1113/jphysiol.2013.270389>
- Fasano A. (2022). *Gut Feelings*. MIT press.

- Fasano A. (2020).** All disease begins in the (leaky) gut: role of zonulin-mediated gut permeability in the pathogenesis of some chronic inflammatory diseases. *F1000Research*, 9, F1000 Faculty Rev-69. <https://doi.org/10.12688/f1000research.20510.1>
- De Filippis F., Valentino V., Sequino G., Borriello G., Riccardi M.G., Pierri B., Cerino P., Pizzolante A., Pasolli E., Esposito M., Limone A., & Ercolini D. (2024).** Exposure to environmental pollutants selects for xenobiotic-degrading functions in the human gut microbiome. *Nature Communications*, 15(1), 4482. <https://doi.org/10.1038/s41467-024-48739-7>
- Foster J.A., & McVey Neufeld K.A. (2013).** Gut-brain axis: how the microbiome influences anxiety and depression. *Trends in Neurosciences*, 36(5), 305–312. <https://doi.org/10.1016/j.tins.2013.01.005>
- Garrett W.S., Lord G.M., Punit S., Lugo-Villarino G., Mazmanian S.K., Ito S., Glickman J.N., & Glimcher L.H. (2007).** Communicable ulcerative colitis induced by T-bet deficiency in the innate immune system. *Cell*, 131(1), 33–45. <https://doi.org/10.1016/j.cell.2007.08.017>
- Gasbarrini A., Dionisi T., & Gasbarrini G. (2019).** L'azione del Microbiota nel trapianto fecale. *Atti della Accademia Lancisiana*, LXIII(1), 113-121.
- Ge L., Liu S., Li S., Yang J., Hu G., Xu C., & Song W. (2022).** Psychological stress in inflammatory bowel disease: Psychoneuroimmunological insights into bidirectional gut-brain communications. *Frontiers in Immunology*, 13, 1016578. <https://doi.org/10.3389/fimmu.2022.1016578>
- Gottlieb G. (2000).** Environmental and behavioral influences on gene activity. *Current Directions in Psychological Science*, 9(3), 93-97. <https://doi.org/10.1111/1467-8721.00068>
- Ichmann-Diounou H., & Menard S. (2020).** Psychological stress, intestinal barrier dysfunctions, and autoimmune disorders: An overview. *Frontiers in Immunology*, 11, 1823. <https://doi.org/10.3389/fimmu.2020.01823>
- Kelly J.R., Borre Y., O'Brien C., Patterson E., El Aidy S., Deane J., Kennedy P.J., Beers S., Scott K., Moloney G., Hoban A.E., Scott L., Fitzgerald P., Ross P., Stanton C., Clarke G., Cryan J.F., & Dinan T.G. (2016).** Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat. *Journal of Psychiatric Research*, 82, 109–118. <https://doi.org/10.1016/j.jpsychires.2016.07.019>
- Kim C.S., Shin G.E., Cheong Y., Shin J.H., Shin D.M., & Chun W.Y. (2022).** Experiencing social exclusion changes gut microbiota composition. *Translational Psychiatry*, 12(1), 254. <https://doi.org/10.1038/s41398-022-02023-8>
- Koenig J.E., Spor A., Scalfone N., Fricker A.D., Stombaugh J., Knight R., Angenent L.T., & Ley R.E. (2011).** Succession of microbial consortia in the developing infant gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America*, 108(Suppl 1), 4578–4585. <https://doi.org/10.1073/pnas.1000081107>
- Kumar H., Lund R., Laiho A., Lundelin K., Ley R.E., Isolauri E., & Salminen S. (2014).** Gut microbiota as an epigenetic regulator: pilot study based on whole-genome methylation analysis. *mBio*, 5(6), e02113-14. <https://doi.org/10.1128/mBio.02113-14>
- Li Q., & Zhou J.M. (2016).** The microbiota-gut-brain axis and its potential therapeutic role in autism spectrum disorder. *Neuroscience*, 324, 131–139. <https://doi.org/10.1016/j.neuroscience.2016.03.013>
- López-Otín C., Blasco M.A., Partridge L., Serrano M., & Kroemer G. (2013).** The hallmarks of aging. *Cell*, 153(6), 1194–1217. <https://doi.org/10.1016/j.cell.2013.05.039>
- Mangiola F., Ianiro G., Franceschi F., Fagioli S., Gasbarrini G., & Gasbarrini A. (2016).** Gut microbiota in autism and mood disorders. *World Journal of Gastroenterology*, 22(1), 361–368. <https://doi.org/10.3748/wjg.v22.i1.361>
- Mohr A.E., Jäger R., Carpenter K.C., Kerksick C.M., Purpura M., Townsend J.R., ... Antonio J. (2020).** The athletic gut microbiota. *Journal of the International Society of Sports Nutrition*, 17(1). <https://doi.org/10.1186/s12970-020-00353-w>
- Mousavi S.E., Delgado-Saborit J.M., Adivi A., Pauwels S., & Godderis L. (2022).** Air pollution and endocrine disruptors induce human microbiome imbalances: A systematic review of recent evidence and possible biological mechanisms. *The Science of the Total Environment*, 816, 151654. <https://doi.org/10.1016/j.scitotenv.2021.151654>
- Neroni B., Evangelisti M., Radocchia G., Di Nardo G., Pantanella F., Villa M.P., & Schippa S. (2021).**

- Relationship between sleep disorders and gut dysbiosis: What affects what? *Sleep Medicine*, 87, 1–7. <https://doi.org/10.1016/j.sleep.2021.08.003>
- Ottaviani E., Ventura N., Mandrioli M., Candela M., Franchini A., & Franceschi C. (2011).** Gut microbiota as a candidate for lifespan extension: An ecological/evolutionary perspective targeted on living organisms as metaorganisms. *Biogerontology*, 12(6), 599–609. <https://doi.org/10.1007/s10522-011-9352-5>
- Ottman N., Smidt H., de Vos W.M., & Belzer C. (2012).** The function of our microbiota: Who is out there and what do they do? *Frontiers in Cellular and Infection Microbiology*, 2, 104. <https://doi.org/10.3389/fcimb.2012.00104>
- Putignani L., & Dallapiccola B. (2016).** Foodomics as part of the host-microbiota-exposome interplay. *Journal of Proteomics*, 147, 3–20. <https://doi.org/10.1016/j.jpro.2016.04.033>
- Rodrigues-Amorim D., Rivera-Baltanás T., Regueiro B., Spuch C., de Las Heras M.E., Vázquez-Noguerol Méndez R., Nieto-Araujo M., Barreiro-Villar C., Olivares J.M., & Agís-Balboa R.C. (2018).** The role of the gut microbiota in schizophrenia: Current and future perspectives. *The World Journal of Biological Psychiatry*, 19(8), 571–585. <https://doi.org/10.1080/15622975.2018.1433878>
- Rothschild D., Weissbrod O., Barkan E., et al. (2018).** Environment dominates over host genetics in shaping human gut microbiota. *Nature*, 555, 210–215. <https://doi.org/10.1038/nature25973>
- Sen P., Molinero-Perez A., O’Riordan K.J., McCafferty C.P., O’Halloran K.D., & Cryan J.F. (2021).** Microbiota and sleep: Awakening the gut feeling. *Trends in Molecular Medicine*, 27(10), 935–945. <https://doi.org/10.1016/j.molmed.2021.07.004>
- Sender R., Fuchs S., & Milo R. (2016a).** Are we really vastly outnumbered? Revisiting the ratio of bacterial to host cells in humans. *Cell*, 164(3), 337–340. <https://doi.org/10.1016/j.cell.2016.01.013>
- Sender R., Fuchs S., & Milo R. (2016b).** Revised estimates for the number of human and bacteria cells in the body. *PLoS Biology*, 14(8), e1002533. <https://doi.org/10.1371/journal.pbio.1002533>
- Sharon G., Cruz N.J., Kang D.W., Gandal M.J., Wang B., Kim Y.M., Zink E.M., Casey C.P., Taylor B.C., Lane C.J., Brammer L.M., Isern N.G., Hoyt D.W., Noecker C., Sweredoski M.J., Moradian A., Borenstein E., Jansson J.K., Knight R., Metz T.O., ... Mazmanian S. K. (2019).** Human gut microbiota from autism spectrum disorder promote behavioral symptoms in mice. *Cell*, 177(6), 1600–1618.e17. <https://doi.org/10.1016/j.cell.2019.05.004>
- Simpson C.A., Diaz-Arteche C., Eliby D., Schwartz O.S., Simmons J.G., & Cowan C. (2021).** The gut microbiota in anxiety and depression – A systematic review. *Clinical Psychology Review*, 83, 101943. <https://doi.org/10.1016/j.cpr.2020.101943>
- Smith R.P., Easson C., Lyle S.M., Kapoor R., Donnelly C.P., Davidson E.J., Parikh E., Lopez J.V., & Tartar J.L. (2019).** Gut microbiome diversity is associated with sleep physiology in humans. *PLoS ONE*, 14(10), e0222394. <https://doi.org/10.1371/journal.pone.0222394>
- Thaiss C.A., Zeevi D., Levy M., Zilberman-Schapira G., Suez J., Tengeler A.C., ... Elinav E. (2014).** Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell*, 159(3), 514–529. <https://doi.org/10.1016/j.cell.2014.09.048>
- Valitutti F., Cucchiara S., & Fasano A. (2019).** Celiac disease and the microbiome. *Nutrients*, 11(10), 2403. <https://doi.org/10.3390/nu11102403>
- Voigt R.M., Forsyth C.B., Green S.J., Engen P.A., & Keshavarzian A. (2016).** Circadian rhythm and the gut microbiome. *International Review of Neurobiology*, 131, 193–205. <https://doi.org/10.1016/bs.irm.2016.07.002>
- Winter G., Hart R.A., Charlesworth R.P.G., & Sharpley C.F. (2018).** Gut microbiome and depression: What we know and what we need to know. *Reviews in the Neurosciences*, 29(6), 629–643. <https://doi.org/10.1515/revneuro-2017-0072>
- Wu W.L., Adame M.D., Liou C.W., et al. (2021).** Microbiota regulate social behaviour via stress response neurons in the brain. *Nature*, 595, 409–414. <https://doi.org/10.1038/s41586-021-03669-y>
- Zhang N., Ju Z., & Zuo T. (2018).** Time for food: The impact of diet on gut microbiota and human health. *Nutrition*, 51-52, 80–85. <https://doi.org/10.1016/j.nut.2017.12.005>